

Conditional escape of gravitons from the brane

Jannie A. Leach ^{†‡} and William M. Lesame [†]

[†] Department of Mathematical Sciences, University of South Africa, Pretoria 0003, South Africa

E-mail: leachja@nassp.uct.ac.za and lesamwm@unisa.ac.za

Abstract. In this paper we consider a cosmological Friedmann Robertson Walker brane world embedded in a 5-dimensional anti-de Sitter Schwarzschild bulk. We show, using potential diagrams, that for an anti-de Sitter bulk the null geodesics never return to the brane. Null geodesics do however return for $k = +1$ when we include the Schwarzschild like mass and the condition of return is obtained from the corresponding effective potential. We next obtain the condition for a gravitational signal to be emitted from a FRW brane with isotropically distributed matter. We use these results to investigate the conditions under which shortcuts through the bulk are possible.

PACS numbers: 98.80.Cq, 04.50.+h

1. Introduction

Causality plays an important role in the brane-world scenario. Null geodesics passing through the extra dimension may connect points in the lower dimension which are causally disconnected in terms of the null geodesics confined to the brane [1]. Thus instead of inflation, such null geodesics could be used to solve the cosmological horizon problem. The condition for the existence of such “causality violating” null geodesics, which pass through the AdS_5 bulk, is merely the deviation from a pure tension-like stress energy tensor, i.e. provided that the energy density ρ_b and the isotropic pressure p_b of the brane satisfies [2]

$$\rho_b + p_b > 0,$$

then the extrinsic curvature bends the brane concave towards the bulk, allowing the existence of such null geodesics.

Initially most authors considered these “shortcuts” through the bulk for a static cosmological brane (brane based approach). They made use of the most general metric (see [3, 4]) of the form

$$ds^2 = -n^2(t, y) dt^2 + a^2(t, y) \gamma_{\mu\nu} dx^\mu dx^\nu + b^2(t, y) dy^2.$$

Null bulk geodesic equations of motion has been computed for static and cosmological single brane models (see [5–7]). It was found that although there are effective acausality in some cases, there is no *a priori* solution to the horizon problem.

[‡] Present address: Department of Mathematics and Applied Mathematics, University of Cape Town, Rondebosch, 7701, South Africa

A more realistic scenario was found by considering a dynamical brane (bulk based approach). It has been shown that the motion of a domain wall in the bulk can be interpreted as cosmological expansion or contraction on the wall (see [8]). A generalization of Birkhoff's theorem can be used to show that a fixed wall embedded in a non-static spacetime is strictly equivalent to a moving wall embedded in a static spacetime and so establishing in full generality the equivalence between the two approaches (see [9]). Solutions have also been found for higher dimensional dynamic domain walls that couples to bulk matter [10].

Causality can now be investigated in a different way. The trajectories of the null geodesics and the brane's equations of motion in the bulk are first obtained. These are then plotted on the same set of axes, with both curves having the same initial conditions. If the curves intersect again, then it can be said that two points are causally connected through the bulk. The brane's equations of motion for a FRW brane in a AdS_5 bulk (only considering the tension of the brane and not the matter inside) was obtained in [8]. Shortcuts through the bulk could thus be investigated from the dynamical or bulk based approach (see e.g. [11, 12]). A FRW brane embedded in a AdS_5 was used and it was found that although the distance covered by a graviton passing through the bulk is longer than the corresponding distance on the brane, it is not enough to solve the horizon problem [11]. Shortcuts have also been investigated in various other models, such as; six dimensional brane worlds [13], $(D - 2)$ -branes (domain walls) [14] and for nonlinear dynamical universes [15].

In none of the references above (and no reference to our knowledge), has the condition of escape been determined for a graviton to actually leave the brane. So far most authors have assumed that gravitons escape freely into the bulk. If this was the case, why then don't all the gravitons escape into the bulk? This is the main question which we will try and answer in this paper. We will attempt to firstly determine whether null geodesics return to the brane after they have escaped and secondly the condition for a graviton to escape from the brane.

2. Null geodesic equations

We start with the 5-dimensional anti-de Sitter Schwarzschild metric

$$ds^2 = -f_k(R)dT^2 + f_k(R)^{-1}dR^2 + R^2 d\Sigma_k^2 \quad (1)$$

where T is the global bulk time, R is the bulk dimension and $d\Sigma_k$ is the metric for a maximally symmetric three-dimensional spaces ($k = 0$ for a flat space, $k = 1$ for a three sphere and $k = -1$ for a hyperbolic three space). The function $f_k(R)$ is given by

$$f_k(R) = k + \frac{R^2}{\ell^2} - \frac{\mu}{R^2} \quad (2)$$

where $\ell \equiv \sqrt{\frac{-6}{\Lambda}}$ is the anti-de Sitter radius of curvature, Λ is the negative bulk cosmological constant (i.e. $\Lambda < 0$) and μ is the five dimensional Schwarzschild-like mass. Since the third term in the equation above scales like radiation, μ is sometimes called the dark radiation or Weyl parameter. The solutions are anti-de Sitter when $\mu = 0$ and when $\mu \neq 0$ we have a bulk black hole with a horizon at

$$R_h^2 = \frac{\ell^2}{2} \left(-k + \sqrt{k^2 + \frac{4\mu}{\ell^2}} \right). \quad (3)$$

A spherical coordinate system (r, θ, ϕ) can now be introduced in the brane which is centered at a point A, so that any path can be described by a radial geodesic [11]. Then due to the radial geodesics we may ignore the angular variables θ and ϕ , so that we are left with a three dimensional problem with the resulting metric being given by

$$ds^2 = -f_k(R)dT^2 + f_k(R)^{-1}dR^2 + R^2dr^2. \quad (4)$$

The geodesics may be easily obtained by using the symmetries to obtain the Killing vectors, namely $(\frac{\partial}{\partial T})^a$ and $(\frac{\partial}{\partial r})^a$ (see [11] and [16]). Then the vector tangent to the geodesics, p^a , is given by

$$p_T = -f_k(R)\frac{dT}{d\lambda} = -E \quad (a \text{ constant}) \quad (5)$$

and

$$p_r = R^2\frac{dr}{d\lambda} = P \quad (a \text{ constant}) \quad (6)$$

where λ is an affine parameter that relates each of the three parameters (R, T, r) to one another. Both E and P are positive constants which are similarly defined as the “energy at infinity” E and “angular momentum” L in standard general relativity (see [16]).

For a null geodesic we furthermore have

$$\left(\frac{dR}{d\lambda}\right)^2 = E^2 - P^2\frac{f_k(R)}{R^2} \quad (7)$$

Equations (5), (6) and (7) may be rewritten in the standard GR form of [16] by setting $\tilde{\lambda} = P\lambda$ so that the full set of geodesic equations are

$$\frac{dT}{d\tilde{\lambda}} = \frac{E}{Pf_k(R)}, \quad (8)$$

$$\frac{dr}{d\tilde{\lambda}} = \frac{1}{R^2}, \quad (9)$$

$$\left(\frac{dR}{d\tilde{\lambda}}\right)^2 = \frac{E^2}{P^2} - \frac{f_k(R)}{R^2}. \quad (10)$$

3. Brane motion

We start by introducing the proper time t on the brane, which is defined by [11] as

$$dt^2 = f(R_b)dT^2 - f(R_b)^{-1}dR_b^2 \quad (11)$$

so that

$$\frac{dT}{dt} = \frac{\sqrt{f(R_b) + \dot{R}_b^2}}{f(R_b)} \quad (12)$$

where $\dot{R}_b = \frac{dR_b}{dt}$. $R_b(t) \equiv a(t)$ is the usual cosmological scale factor in the brane-world which can be obtained from the modified Friedmann equation which is determined from the junction conditions.

The Israel junction conditions [17] relate the discontinuity in the second fundamental form (extrinsic curvature) through the brane and the energy momentum

tensor in the brane. If K_{ab}^+ and K_{ab}^- are the jump in extrinsic curvature on the two sides of the brane, then the junction conditions for a Z_2 symmetric brane is [2]

$$K_{ij}^+ = -K_{ij}^- = -\frac{1}{2}\kappa_5^2 \left(S_{ij} - \frac{1}{3}h_{ij}S \right). \quad (13)$$

The extrinsic curvature of the hypersurface can be computed using the unit norm vector (see e.g. [12, 18, 19]). If we assume an isotropic distribution of matter on the brane, then the energy momentum tensor S_{ij} in the brane is given by

$$S_{ij} = \rho_b u_i u_j + p_b (h_{ij} + u_i u_j), \quad (14)$$

where ρ_b is the energy density and p_b is the isotropic pressure of the brane.

From the junction conditions above one obtains two sets of equation governing the evolution of the brane (see [3, 8, 9]):

$$\frac{d\rho_b}{dt} = -3\frac{\dot{R}_b}{R_b}(\rho_b + p_b). \quad (15)$$

and

$$\frac{\dot{R}_b^2}{R_b^2} + \frac{f(R_b)}{R_b^2} = \frac{\kappa_5^4 \rho_b^2}{36}. \quad (16)$$

We require some equation of state that relates the matter density ρ and pressure p on the brane. We follow convention by setting

$$\rho_b = \rho + \sigma, \quad p_b = p - \sigma,$$

where σ is tension of the wall. The equation of state then may be given by

$$p = (\gamma - 1)\rho.$$

Substitution then yields the conservation equations as

$$\frac{d\rho}{dt} = -3\frac{\dot{R}_b}{R_b}(\rho + p), \quad (17)$$

with the solution

$$\rho = \rho_0 \left(\frac{R_0}{R_b} \right)^{3\gamma}$$

where the subscript 0 means any given time. Similarly we may obtain the modified Friedmann equation of [4] in the form

$$\dot{R}_b^2 + \left(1 - \left(\frac{\rho + \sigma}{\sigma_c} \right)^2 \right) \frac{R_b^2}{\ell^2} - \frac{\mu}{R_b^2} = -k, \quad (18)$$

where $\sigma_c \equiv 3/(4\pi G\ell^2) = 6/(\kappa_5^2 \ell)$ is the critical tension for fine tuning. The cosmological constant in the brane then has the form

$$\frac{\Lambda_4}{3} = \frac{\kappa_5^4 \sigma^2}{36} - \frac{1}{\ell^2} = \frac{1}{\ell^2} \left(\frac{\sigma^2}{\sigma_c^2} - 1 \right).$$

4. Graviton emission and capture

4.1. Potential diagrams

In the case of a domain wall, there is no matter on the wall so we do not expect the escaping gravitons to be attracted back to the brane. A more realistic scenario would be to consider a brane with isotropically distributed matter. Analytically computing trajectories of the moving brane with matter is difficult so we make use of potential diagrams to study whether gravitational signals will return to the brane. In classical GR, potential diagrams have been found useful for qualitative analysis of orbits of particles and photons in Schwarzschild geometry. Potential diagrams has been used before to investigate Lorentz violations in $AdSS_5$ bulk background [20]. Here we extend that analysis to brane motion and graviton motion in AdS_5 and $AdSS_5$.

We start by rewriting equation (10) as

$$\left(\frac{dR}{d\lambda}\right)^2 + B^2(R) = b^2, \quad (19)$$

where

$$B^2(R) = \frac{f_k(R)}{R^2} \quad (20)$$

defines a kind of “effective potential of a graviton”, and

$$b^2 = \frac{E^2}{P^2} \quad (21)$$

defines the “impact parameter”.

The potential impact parameter $B(R)$ has the following useful interpretation: For a graviton to reach the point R , it must have a impact parameter b greater than or equal to $B(R)$, i.e.

$$b \geq B(R). \quad (22)$$

This is referred to as the *condition of accessibility* [16].

AdS_5 ($\mu = 0$): The effective potential (20) for the various values of k are

$$B^2(R) = \frac{1}{\ell^2} + \frac{1}{R^2}, \quad k = +1 \quad (23)$$

$$B^2(R) = \frac{1}{\ell^2}, \quad k = 0 \quad (24)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{1}{R^2}, \quad k = -1 \quad (25)$$

and are plotted in Figure 1. Clearly none of the potential curves have turning points, i.e if a graviton escape from the brane (domain wall or brane with matter) into an anti-de Sitter bulk, it will not return to the brane. For $k = -1$ we have a horizon at $R = \ell$.

$AdSS_5$ ($\mu \neq 0$): The effective potential (20) for the various values of k are

$$B^2(R) = \frac{1}{\ell^2} + \frac{1}{R^2} - \frac{\mu}{R^4}, \quad k = +1 \quad (26)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{\mu}{R^4}, \quad k = 0 \quad (27)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{1}{R^2} - \frac{\mu}{R^4}, \quad k = -1 \quad (28)$$




Figure 1. Potential diagram for AdS_5 ($\mu = 0$) bulk geometry.




Figure 2. Potential diagram for $\mu \neq 0$ and $k = +1$.

The case where $k = +1$ is plotted in Figure 2. Both the $k = 0$ and $k = -1$ cases look similar to $k = -1$ when $\mu = 0$ (see Figure 1), i.e they have no turning points. The horizons for all k 's are given by (3).

4.2. Condition of return

The turning point on Figure 2, is at

$$R_t = (2\mu)^{\frac{1}{2}} \quad (29)$$

which implies a maximum effective potential of

$$B^2(R_t) = \frac{1}{\ell^2} + \frac{1}{4\mu}.$$

Combining this with the condition of accessibility (22), we can summarize the qualitative features of the trajectories of gravitons emitted in the $AdSS_5$ bulk with $k = +1$ (see [16]) as follows:

When a graviton is emitted near the horizon R_h (i.e. at some point R_{em} , $R_h < R_{em} < R_t$), it will:

(i) escape to infinity when

$$b > \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}; \quad (30)$$

(ii) or return when

$$0 < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}. \quad (31)$$

Equation (31) in effect gives us the *condition of return* for a graviton emitted near the bulk horizon R_h . The lower limit arise from the fact that both E and P are positive constants, and hence $b > 0$. Clearly we can only expect returning gravitons at early times when $R_{em} < R_t$ when the effective potential is increasing. When a graviton is emitted at $R_{em} > R_t$ it will escape to infinity (since $B(R)$ is decreasing).

4.3. Conditions of escape

For a gravitational signal to escape from the brane, it must move faster than the brane with respect to an observer in the bulk. Thus the relative velocity of the gravitational signal v_g must be greater than the relative velocity of the brane v_b at the point of emission R_{em} . Therefore, we now seek to obtain the relative velocities of the gravitational signal and the brane.

We first obtain the relative velocity of the gravitational signal by dividing equation (8) into equation (10), and then taking the square root to find

$$v_g = \frac{dR_g}{dT} = f_k(R_g) \sqrt{1 - \frac{P^2 f_k(R_g)}{E^2 R_g^2}} = f_k(R_g) \sqrt{1 - \frac{B^2(R_g)}{b^2}}. \quad (32)$$

We see that this equation will hold if the condition of accessibility (22) is satisfied (where we made use of (20) and (21)). The velocity will be zero when $b = B(R_g)$, i.e. we will have a turning point for a gravitational signal.

Next we find the relative velocity of the brane by substituting equation (11) for dt into (16),

$$dR_b^2 = \left[\left(\frac{\rho + \sigma}{\sigma_c} \right)^2 \frac{R_b^2}{\ell^2} - f_k(R_b) \right] [f_k(R_b) dT^2 - f_k^{-1}(R_b) dR_b^2], \quad (33)$$

which on simplification yields

$$v_b = \frac{dR_b}{dT} = f_k(R_b) \sqrt{1 - \frac{\ell^2 f_k(R_b)}{\left(\frac{\rho + \sigma}{\sigma_c} \right)^2 R_b^2}}. \quad (34)$$

The equation for the relative velocity of the brane will only hold if

$$\hat{\beta} = \frac{1}{\ell} \left(\frac{\rho + \sigma}{\sigma_c} \right) \geq B(R_b). \quad (35)$$

In the limit of $R \rightarrow 0$ this imply that $\mu \geq 0$ and in the limit of $R \rightarrow \infty$ we must have $\sigma \geq \sigma_c$. We are therefore only considering positive dark radiation.

The ratio of the two relative velocities is then given by

$$\frac{v_g}{v_b} = \frac{f_k(R_g)}{f_k(R_b)} \cdot \sqrt{\frac{1 - \frac{P^2 f_k(R_g)}{E^2 R_g^2}}{1 - \frac{\ell^2 f_k(R_b)}{\left(\frac{\rho + \sigma}{\sigma_c}\right)^2 R_b^2}}}. \quad (36)$$

We seek to find the conditions of escape, and therefore we only consider the position of the brane and the energy density at the time of emission, i.e. $R_b = R_g = R_{em}$ and $\rho = \rho_{em}$ respectively. The graviton will have the same position as the brane at the bulk time of emission. A gravitational signal can be emitted if and only if $v_g > v_b$, that is when

$$1 - \frac{P^2 f_k(R_{em})}{E^2 R_{em}^2} > 1 - \frac{\ell^2 f_k(R_{em})}{\left(\frac{\rho_{em} + \sigma}{\sigma_c}\right)^2 R_{em}^2},$$

which after simplification reduces to

$$\frac{E}{P} > \frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) \quad (37)$$

or

$$b > \hat{\beta}_{em}. \quad (38)$$

This condition can be considered as the *condition of escape*, that is the condition which needs to be satisfied for a graviton to escape from the brane.

When $v_g = v_b$ we have the condition

$$\frac{E}{P} = \frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) \quad (39)$$

which implies that the graviton is stuck on the brane and therefore appears to be moving with the brane at the same relative velocity. In the case when $v_g < v_b$, the gravitational signal is lagging the brane. This is clearly not possible.

5. Summary and discussion

The only case which showed the possibility of emitting a graviton with it returning to the brane, is $k = +1$ when $\mu \neq 0$. The potential diagram for this case has a classical potential form (see [16] and Figure 2) which meant that if a graviton is emitted from the brane at R_{em} (where $R_h < R_{em} < R_t$) it would return if the condition (31) is satisfied. Comparing the velocities of the graviton and the brane we were able to determine that a graviton will be able to escape from the brane if the condition (37) is satisfied.

The condition for a graviton to escape from the brane at some early time (when $R_h < R_{em} < R_t$) and return, is then given by

$$\frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}, \quad (40)$$

where $\mu > 0$ and $\sigma \geq \sigma_c$. This condition can also be expressed in terms of the 5-dimensional fundamental mass scale M_5 , by using the relationship $\kappa_5^2 = 8\pi G\ell = M_5^{-3}$, namely

$$\frac{\rho_{em} + \sigma}{6M_5^3} < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}. \quad (41)$$

Since the impact parameter has no physical significance, we rather use the conditions above to find the relationship between parameters that do have physical significance. We find that the conditions of escape and return impose the following restrictions on the dark radiation μ (Schwarzschild-like mass)

$$\mu < \frac{\ell^2}{4} \left[\left(\frac{\rho_{em} + \sigma}{\sigma_c} \right)^2 - 1 \right]^{-1}, \quad (42)$$

for all $\rho_{em} + \sigma > \sigma_c$. This in turn tells us that we may only have returning gravitons before the brane reaches the point R_t , where using (29) we have the condition

$$R_t < \frac{\ell}{\sqrt{2}} \left[\left(\frac{\rho_{em} + \sigma}{\sigma_c} \right)^2 - 1 \right]^{-\frac{1}{2}}. \quad (43)$$

We can realistically only expect emitted gravitons to return to the brane at very early times ($\rho_{em} + \sigma \gg \sigma_c$) when $R_{em} \rightarrow R_h$ and thus $B(R_{em}) \rightarrow 0$. At late times ($\rho \rightarrow 0$) we are essentially looking at a domain wall moving in an AdS_5 space where gravitons do not return to the brane (see appendix). In the limit $\rho_{em} + \sigma \gg \sigma_c$ the restrictions above reduce to

$$\mu < \frac{1}{(\rho_{em} + \sigma)^2} \quad (44)$$

and hence

$$R_t < \frac{1}{\rho_{em} + \sigma}. \quad (45)$$

The upper limit for the dark radiation μ obtained here still require further investigation, especially in the light of the nucleosynthesis limits [21].

We see that there exist a small window of opportunity, before the brane reaches R_t , for gravitons to be emitted from the brane and then return to it. After the brane reaches R_t , all gravitons emitted never return to the brane. An alternative might be to look at a bulk with some sort of field (dilaton etc.) which forces the path back to the brane when it has gone beyond R_t . Another possibility is to have a black hole on the brane that will give us the required curvature. Black holes on the brane have been consider by a number of authors (see [22–26]) most notably in the context of causality, by [27].

Another issue that needs investigation, is that of conservation of energy for a brane that is radiating these graviton. Clearly the form (15) does not take into consideration the gravitons that are emitted. Radiating braneworlds have been considered in [28] and [29], which might show more promising results. The mechanisms that could produce gravitons with sufficient energy to leave the brane has also as yet not been explored properly.

In conclusion, there are two important aspect of escaping gravitational signals. In §4 we clearly saw that the condition of return does not depend on the type of brane that we choose, but rather on the type of bulk (i.e. on its 5-D spacetime). We

also discovered that the condition of escape depends on the type of brane we choose and is independent of the bulk space time. We note that no matter which values of μ (AdS_5) and charge Q (Reissner-Nordström) we choose, the escape condition still stays the same. The condition of escape is also independent of the curvature values k .

We have thus shown that there are certain conditions which first needs to be satisfied before one may assume that a signal can escape and then return to the brane. Two important conditions need to be satisfied, firstly we need a condition under which a graviton can escape from the brane (or be captured by the brane), and secondly the conditions which allow it to return to the brane. Furthermore, these two conditions together restrict gravitons to escaping and returning to the brane before the brane reaches a point R_t in the bulk. If all of these conditions are not satisfied one can not expect to have causally connected points through the bulk.

Acknowledgments

This research was supported by the National Research Foundation (South Africa). J.L. also acknowledges the postgraduate assistantship from the University of South Africa (UNISA).

Appendix A. Domain wall motion in AdS_5

In this section we give the results for the special case where we are considering a empty brane (domain wall) moving in an AdS_5 bulk.

Appendix A.1. Null geodesics solutions for AdS_5

When the bulk is anti-de Sitter AdS_5 , i.e $\mu = 0$, one can readily obtain the solutions of the null geodesic equations (8)-(10). For $k = 0$ the solutions are

$$R_g = \frac{\mathcal{B}_g}{\mathcal{A}_g - T}, \quad \alpha > 0, \quad (\text{A.1})$$

and

$$R_g = \frac{\mathcal{B}_g}{T - \mathcal{A}_g}, \quad \alpha < 0, \quad (\text{A.2})$$

where $\alpha^2 = \frac{E^2}{P^2} - \frac{1}{\ell^2}$ and

$$\mathcal{A}_g = T_0 \pm \frac{E\ell^2}{\alpha P R_0}, \quad \mathcal{B}_g = \frac{E\ell^2}{\alpha P}.$$

The + sign in $\mathcal{A}_g$ holds when $\alpha > 0$, the - sign when $\alpha < 0$.

In the case where $k = -1$, the solutions are

$$R_g^2 = \frac{E^2\ell^2}{\alpha^2 P^2} \coth^2 \left(\frac{\mathcal{C}_g - T}{\ell} \right) - \frac{1}{\alpha^2}, \quad R > \ell, \quad (\text{A.3})$$

and

$$R_g^2 = \frac{E^2\ell^2}{\alpha^2 P^2} \tanh^2 \left(\frac{\mathcal{D}_g - T}{\ell} \right) - \frac{1}{\alpha^2}, \quad R < \ell, \quad (\text{A.4})$$

where the initial conditions are

$$\mathcal{C}_g = T_0 + \ell \coth^{-1} \frac{P\sqrt{\alpha^2 R_0^2 + 1}}{E\ell},$$

and

$$\mathcal{D}_g = T_0 + \ell \tanh^{-1} \frac{P\sqrt{\alpha^2 R_0^2 + 1}}{E\ell}.$$

Similarly for $k = +1$ the solutions are

$$R_g^2 = \frac{E^2 \ell^2}{\alpha^2 P^2} \tan^2 \left(\frac{T - \mathcal{F}_g}{\ell} \right) + \frac{1}{\alpha^2}, \quad (\text{A.5})$$

where

$$\mathcal{F}_g = T_0 - \ell \tan^{-1} \frac{P\sqrt{\alpha^2 R_0^2 - 1}}{E\ell}.$$

The solutions to the geodesic equation for the radial coordinate r in terms of the global bulk time T , for all $k = 0, -1$ and $+1$, are given by

$$r = r_0 + \frac{P}{E\ell^2} (T - T_0). \quad (\text{A.6})$$

Appendix A.2. Trajectories for a moving domain wall

When one considers a domain wall (i.e. $\rho = 0$) moving in an anti-de Sitter bulk ($\mu = 0$), the modified Friedmann equation (18) takes the form [8]

$$\dot{R}_b^2 + \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \right) \frac{R_b^2}{\ell^2} = -k. \quad (\text{A.7})$$

The solutions of these, in terms of the proper time on the brane t , are given in Kraus [8]. We are however interested in the brane's trajectory in the bulk. We therefore transform Kraus' equations by making use of (12). The trajectories are

$k = 0, \sigma = \sigma_c$: $R_b = \text{constant}$

$k = 0, \sigma > \sigma_c$:

solution 1 (expanding brane):

$$R_b = \frac{1}{H} \left(\frac{\sigma}{\sigma_c} \right) \frac{\ell}{\mathcal{A}_b - T}, \text{ where } \mathcal{A}_b = T_0 + \frac{\sigma}{\sigma_c} \frac{\ell}{R_0 H}, \quad (\text{A.8})$$

and where $H = \frac{1}{\ell} \sqrt{\left(\frac{\sigma}{\sigma_c} \right)^2 - 1}$.

solution 2 (contracting brane):

$$R_b = \frac{1}{H} \left(\frac{\sigma}{\sigma_c} \right) \frac{\ell}{T - \mathcal{A}_b}, \text{ where } \mathcal{A}_b = T_0 - \frac{\sigma}{\sigma_c} \frac{\ell}{R_0 H}. \quad (\text{A.9})$$

$k = -1, \sigma = \sigma_c$:

$$R_b^2 = \ell^2 - (\ell^2 - R_0^2) e^{\frac{2(T-T_0)}{\ell}}, \quad R < \ell, \quad (\text{A.10})$$

and

$$R_b^2 = (R_0^2 - \ell^2) e^{\frac{2(T-T_0)}{\ell}} + \ell^2, \quad R > \ell. \quad (\text{A.11})$$

$k = -1, \sigma > \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \coth^2 \left(\frac{\mathcal{C}_b - T}{\ell} \right) - 1 \right), \quad R > \ell, \quad (\text{A.12})$$

and

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \tanh^2 \left(\frac{\mathcal{D}_b - T}{\ell} \right) - 1 \right), \quad R < \ell, \quad (\text{A.13})$$

where

$$\mathcal{C}_b = T_0 + \ell \coth^{-1} \left(\frac{\sqrt{H^2 R_0^2 + 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right),$$

and

$$\mathcal{D}_b = T_0 + \ell \tanh^{-1} \left(\frac{\sqrt{H^2 R_0^2 + 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$

$k = -1, \sigma < \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \coth^2 \left(\frac{T - \mathcal{C}_{b2}}{\ell} \right) \right), \quad R < \ell, \quad (\text{A.14})$$

and

$$R_b^2 = \frac{1}{H^2} \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \tanh^2 \left(\frac{T - \mathcal{D}_{b2}}{\ell} \right) \right), \quad R > \ell, \quad (\text{A.15})$$

where

$$\mathcal{C}_{b2} = T_0 - \ell \coth^{-1} \left(\frac{\sqrt{1 - H^2 R_0^2}}{\left(\frac{\sigma}{\sigma_c} \right)} \right),$$

and

$$\mathcal{D}_{b2} = T_0 - \ell \tanh^{-1} \left(\frac{\sqrt{1 - H^2 R_0^2}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$

$k = +1, \sigma > \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \tan^2 \left(\frac{T - \mathcal{F}_b}{\ell} \right) + 1 \right), \quad (\text{A.16})$$

where

$$\mathcal{F}_b = T_0 - \ell \tan^{-1} \left(\frac{\sqrt{H^2 R_0^2 - 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$




Figure A1. Expanding brane: R versus T for $k = 0$, $\sigma > \sigma_c$ and $\alpha > 0$.

Appendix A.3. Analysis

We now have the trajectories for both the empty brane (domain wall) and an gravitational signal in AdS_5 . We plot them on the same set of axes, with the same initial conditions so as to see the behaviour of the gravitational signal after it has left the brane.

In order for us to compare the trajectories we set

$$\omega = \frac{E\ell}{P} \quad (\text{A.17})$$

in the null geodesic equations, and set

$$\beta = \frac{\sigma}{\sigma_c} \quad (\text{A.18})$$

in the brane's equations of motion. We assume that $\ell \sim 10^{-4}m$.

In Figure A1 we start with the $k = 0$ case for an expanding universe (solution 1) with $\sigma > \sigma_c$. We set $\beta = 2$ and $R_0 = 1$ at $T_0 = 0$, in (A.8) for the brane equation of motion. For the gravitational signals path (null geodesic), we consider $\omega = 4$ and $\omega = 1.5$ in (A.1) with the same initial conditions. In Figure A2 we plot the results for a contracting universe (solution 2). The two cases for $k = -1$ are plotted in Figure A3 and Figure A4 and the case when $k = 1$ is plotted in Figure A5 (all with $\sigma > \sigma_c$).

Appendix A.3.1. $\omega = \beta$ When we set $\omega = \beta$ we find that the gravitational signal follow the same path as the brane for all cases of k when $\sigma > \sigma_c$. We can interpreted this condition as the graviton being confined to the brane and therefore having the same coordinate in the bulk. We can restate this condition in terms of the impact parameter

$$b = \frac{E}{P} = \frac{1}{\ell} \left(\frac{\sigma}{\sigma_c} \right) = \frac{\sigma}{6M_5^3}. \quad (\text{A.19})$$




Figure A2. Contracting brane: R versus T for $k = 0$, $\sigma > \sigma_c$ and $\alpha < 0$.




Figure A3. R versus T for $k = -1$ and $\sigma > \sigma_c$ and $R > \ell$




Figure A4. R versus T for $k = -1$, $\sigma > \sigma_c$ and $R < \ell$.




Figure A5. R versus T for $k = 1$ and $\sigma > \sigma_c$

Appendix A.3.2. $\omega > \beta$ When $\omega > \beta$ the geodesic equation is not the same as the equation of motion. Thus a graviton can be emitted along the extra-dimensional geodesic, i.e. can be emitted from the brane. This can therefore be seen as the condition for a graviton to be emitted from a domain wall with tension σ into a AdS_5 space. We can express this condition in the form

$$b = \frac{E}{P} > \frac{1}{\ell} \left(\frac{\sigma}{\sigma_c} \right). \quad (\text{A.20})$$

This condition amounts to the gravitational signal having a greater velocity than the brane and may serve as a late time approximation of (37) since at late times the AdS_5 becomes AdS_5 and the brane a domain wall. We can therefore see from Figure A5 that no gravitational signals return to the brane at late times.

Appendix A.3.3. $\omega < \beta$ In the $k = 0$ and $k = -1$ cases we see that the possibility of $\omega < \beta$ also gives paths that are different to the brane's path. However, in these cases the brane is moving faster than the gravitational signal. Clearly these cases are not valid. The initial conditions for the $k = 1$, exclude the possibility of $\omega < \beta$.

References

- [1] Chung D and Freese K, *Phys. Rev.* **D62**, 063513 (2000).
- [2] Ishihara H, *Phys. Rev. Lett.* **86**, 381 (2001).
- [3] Binétruy P, Deffayet C and Langlois D, *Nuc. Phys.* **B565**, 269 (2000).
- [4] Binétruy P, Deffayet C, Ellwanger U and Langlois D, *Phys. Lett.* **B477**, 285 (2000).
- [5] Youm D, *Mod. Phys. Lett.* **A16**, 2371 (2001).
- [6] Davis A, Rhodes C and Vernon I, *JHEP* **0111**, 015 (2001).
- [7] Abdalla E, Cuadros-Melgar B, Feng S and Wang B, *Phys. Rev.* **D65**, 083512 (2002).
- [8] Kraus P, *JHEP* **9912**, 011 (1999).
- [9] Bowcock P, Charmousis C and Gregory R, *Class. Quant. Grav.* **17**, 4745 (2000).
- [10] Chamblin A and Reall H S, *Nuc. Phys.* **B562**, 133 (1999).
- [11] Caldwell R and Langlois D, *Phys. Lett.* **B511**, 129 (2001).
- [12] Abdalla E, Casali A G and Cuadros-Melgar B, *Int. J. Theor. Phys.* **43**, 801 (2004).
- [13] Abdalla E, Casali A G and Cuadros-Melgar B, *Nuc. Phys.* **B644**, 201 (2002).
- [14] Abdalla E and Cuadros-Melgar B, *Phys. Rev.* **D67**, 084012 (2003).
- [15] Cuadros-Melgar B, *Class. Quant. Grav.* **21**, 2669 (2004).
- [16] Misner C W, Thorne K S and Wheeler J A, *Gravitation* (Freeman, San Francisco 1973).
- [17] Israel W, *Nuovo Cimento* **B44**, 1 (1966), erratum: **B48**, 463 (1967).
- [18] Brax P, Langlois D and Rodríguez-Martínez M, *Phys. Rev.* **D67**, 104022 (2003).
- [19] Brax P and van de Bruck C, *Class. Quant. Grav.* **20**, R201 (2003) [arXiv: hep-th/0303095].
- [20] Csáki C, Erlich J and Grojean C, *Nuc. Phys.* **B604**, 312 (2001).
- [21] Ichiki K, Yahiro M, Kajino T, Orito M and Mathews G J, *Phys. Rev.* **D66**, 043521 (2002).
- [22] Chamblin A, Hawking S W and Reall H S, *Phys. Rev.* **D61**, 065007 (2000).
- [23] Dadhich N, Maartens R, Papadopoulos P and Rezanian V, *Phys. Lett.* **B487**, 1 (2000).
- [24] Emparan R, Horowitz G T and Myers R C, *JHEP* **0001**, 007 (2000).
- [25] Chamblin A, Reall H.S, Shinkau H. and Shiromizu T., *Phys. Rev.* **D63**, 064015 (2001).
- [26] Pravda V and Pravdova A, *Class. Quant. Grav.* **18**, 1205 (2001).
- [27] Chamblin A, *Class. Quant. Grav.* **18**, L17 (2001).
- [28] Langlois D, Sorbo L and Rodríguez-Martínez M, *Phys. Rev. Lett.* **89**, 171301 (2002).
- [29] Leeper E, Maartens R and Sopuerta C F, *Class. Quant. Grav.* **21**, 1125 (2004).

Conditional escape of gravitons from the brane

Jannie A. Leach ^{†‡} and William M. Lesame [†]

[†] Department of Mathematical Sciences, University of South Africa, Pretoria 0003, South Africa

E-mail: leachja@nassp.uct.ac.za and lesamwm@unisa.ac.za

Abstract. In this paper we consider a cosmological Friedmann Robertson Walker brane world embedded in a 5-dimensional anti-de Sitter Schwarzschild bulk. We show, using potential diagrams, that for an anti-de Sitter bulk the null geodesics never return to the brane. Null geodesics do however return for $k = +1$ when we include the Schwarzschild like mass and the condition of return is obtained from the corresponding effective potential. We next obtain the condition for a gravitational signal to be emitted from a FRW brane with isotropically distributed matter. We use these results to investigate the conditions under which shortcuts through the bulk are possible.

PACS numbers: 98.80.Cq, 04.50.+h

1. Introduction

Causality plays an important role in the brane-world scenario. Null geodesics passing through the extra dimension may connect points in the lower dimension which are causally disconnected in terms of the null geodesics confined to the brane [1]. Thus instead of inflation, such null geodesics could be used to solve the cosmological horizon problem. The condition for the existence of such “causality violating” null geodesics, which pass through the AdS_5 bulk, is merely the deviation from a pure tension-like stress energy tensor, i.e. provided that the energy density ρ_b and the isotropic pressure p_b of the brane satisfies [2]

$$\rho_b + p_b > 0,$$

then the extrinsic curvature bends the brane concave towards the bulk, allowing the existence of such null geodesics.

Initially most authors considered these “shortcuts” through the bulk for a static cosmological brane (brane based approach). They made use of the most general metric (see [3, 4]) of the form

$$ds^2 = -n^2(t, y) dt^2 + a^2(t, y) \gamma_{\mu\nu} dx^\mu dx^\nu + b^2(t, y) dy^2.$$

Null bulk geodesic equations of motion has been computed for static and cosmological single brane models (see [5–7]). It was found that although there are effective acausality in some cases, there is no *a priori* solution to the horizon problem.

[‡] Present address: Department of Mathematics and Applied Mathematics, University of Cape Town, Rondebosch, 7701, South Africa

A more realistic scenario was found by considering a dynamical brane (bulk based approach). It has been shown that the motion of a domain wall in the bulk can be interpreted as cosmological expansion or contraction on the wall (see [8]). A generalization of Birkhoff's theorem can be used to show that a fixed wall embedded in a non-static spacetime is strictly equivalent to a moving wall embedded in a static spacetime and so establishing in full generality the equivalence between the two approaches (see [9]). Solutions have also been found for higher dimensional dynamic domain walls that couples to bulk matter [10].

Causality can now be investigated in a different way. The trajectories of the null geodesics and the brane's equations of motion in the bulk are first obtained. These are then plotted on the same set of axes, with both curves having the same initial conditions. If the curves intersect again, then it can be said that two points are causally connected through the bulk. The brane's equations of motion for a FRW brane in a AdS_5 bulk (only considering the tension of the brane and not the matter inside) was obtained in [8]. Shortcuts through the bulk could thus be investigated from the dynamical or bulk based approach (see e.g. [11, 12]). A FRW brane embedded in a AdS_5 was used and it was found that although the distance covered by a graviton passing through the bulk is longer than the corresponding distance on the brane, it is not enough to solve the horizon problem [11]. Shortcuts have also been investigated in various other models, such as; six dimensional brane worlds [13], $(D - 2)$ -branes (domain walls) [14] and for nonlinear dynamical universes [15].

In none of the references above (and no reference to our knowledge), has the condition of escape been determined for a graviton to actually leave the brane. So far most authors have assumed that gravitons escape freely into the bulk. If this was the case, why then don't all the gravitons escape into the bulk? This is the main question which we will try and answer in this paper. We will attempt to firstly determine whether null geodesics return to the brane after they have escaped and secondly the condition for a graviton to escape from the brane.

2. Null geodesic equations

We start with the 5-dimensional anti-de Sitter Schwarzschild metric

$$ds^2 = -f_k(R)dT^2 + f_k(R)^{-1}dR^2 + R^2 d\Sigma_k^2 \quad (1)$$

where T is the global bulk time, R is the bulk dimension and $d\Sigma_k$ is the metric for a maximally symmetric three-dimensional spaces ($k = 0$ for a flat space, $k = 1$ for a three sphere and $k = -1$ for a hyperbolic three space). The function $f_k(R)$ is given by

$$f_k(R) = k + \frac{R^2}{\ell^2} - \frac{\mu}{R^2} \quad (2)$$

where $\ell \equiv \sqrt{\frac{-6}{\Lambda}}$ is the anti-de Sitter radius of curvature, Λ is the negative bulk cosmological constant (i.e. $\Lambda < 0$) and μ is the five dimensional Schwarzschild-like mass. Since the third term in the equation above scales like radiation, μ is sometimes called the dark radiation or Weyl parameter. The solutions are anti-de Sitter when $\mu = 0$ and when $\mu \neq 0$ we have a bulk black hole with a horizon at

$$R_h^2 = \frac{\ell^2}{2} \left(-k + \sqrt{k^2 + \frac{4\mu}{\ell^2}} \right). \quad (3)$$

A spherical coordinate system (r, θ, ϕ) can now be introduced in the brane which is centered at a point A, so that any path can be described by a radial geodesic [11]. Then due to the radial geodesics we may ignore the angular variables θ and ϕ , so that we are left with a three dimensional problem with the resulting metric being given by

$$ds^2 = -f_k(R)dT^2 + f_k(R)^{-1}dR^2 + R^2dr^2. \quad (4)$$

The geodesics may be easily obtained by using the symmetries to obtain the Killing vectors, namely $(\frac{\partial}{\partial T})^a$ and $(\frac{\partial}{\partial r})^a$ (see [11] and [16]). Then the vector tangent to the geodesics, p^a , is given by

$$p_T = -f_k(R)\frac{dT}{d\lambda} = -E \quad (a \text{ constant}) \quad (5)$$

and

$$p_r = R^2\frac{dr}{d\lambda} = P \quad (a \text{ constant}) \quad (6)$$

where λ is an affine parameter that relates each of the three parameters (R, T, r) to one another. Both E and P are positive constants which are similarly defined as the “energy at infinity” E and “angular momentum” L in standard general relativity (see [16]).

For a null geodesic we furthermore have

$$\left(\frac{dR}{d\lambda}\right)^2 = E^2 - P^2\frac{f_k(R)}{R^2} \quad (7)$$

Equations (5), (6) and (7) may be rewritten in the standard GR form of [16] by setting $\tilde{\lambda} = P\lambda$ so that the full set of geodesic equations are

$$\frac{dT}{d\tilde{\lambda}} = \frac{E}{Pf_k(R)}, \quad (8)$$

$$\frac{dr}{d\tilde{\lambda}} = \frac{1}{R^2}, \quad (9)$$

$$\left(\frac{dR}{d\tilde{\lambda}}\right)^2 = \frac{E^2}{P^2} - \frac{f_k(R)}{R^2}. \quad (10)$$

3. Brane motion

We start by introducing the proper time t on the brane, which is defined by [11] as

$$dt^2 = f(R_b)dT^2 - f(R_b)^{-1}dR_b^2 \quad (11)$$

so that

$$\frac{dT}{dt} = \frac{\sqrt{f(R_b) + \dot{R}_b^2}}{f(R_b)} \quad (12)$$

where $\dot{R}_b = \frac{dR_b}{dt}$. $R_b(t) \equiv a(t)$ is the usual cosmological scale factor in the brane-world which can be obtained from the modified Friedmann equation which is determined from the junction conditions.

The Israel junction conditions [17] relate the discontinuity in the second fundamental form (extrinsic curvature) through the brane and the energy momentum

tensor in the brane. If K_{ab}^+ and K_{ab}^- are the jump in extrinsic curvature on the two sides of the brane, then the junction conditions for a Z_2 symmetric brane is [2]

$$K_{ij}^+ = -K_{ij}^- = -\frac{1}{2}\kappa_5^2 \left(S_{ij} - \frac{1}{3}h_{ij}S \right). \quad (13)$$

The extrinsic curvature of the hypersurface can be computed using the unit norm vector (see e.g. [12, 18, 19]). If we assume an isotropic distribution of matter on the brane, then the energy momentum tensor S_{ij} in the brane is given by

$$S_{ij} = \rho_b u_i u_j + p_b (h_{ij} + u_i u_j), \quad (14)$$

where ρ_b is the energy density and p_b is the isotropic pressure of the brane.

From the junction conditions above one obtains two sets of equation governing the evolution of the brane (see [3, 8, 9]):

$$\frac{d\rho_b}{dt} = -3\frac{\dot{R}_b}{R_b}(\rho_b + p_b). \quad (15)$$

and

$$\frac{\dot{R}_b^2}{R_b^2} + \frac{f(R_b)}{R_b^2} = \frac{\kappa_5^4 \rho_b^2}{36}. \quad (16)$$

We require some equation of state that relates the matter density ρ and pressure p on the brane. We follow convention by setting

$$\rho_b = \rho + \sigma, \quad p_b = p - \sigma,$$

where σ is tension of the wall. The equation of state then may be given by

$$p = (\gamma - 1)\rho.$$

Substitution then yields the conservation equations as

$$\frac{d\rho}{dt} = -3\frac{\dot{R}_b}{R_b}(\rho + p), \quad (17)$$

with the solution

$$\rho = \rho_0 \left(\frac{R_0}{R_b} \right)^{3\gamma}$$

where the subscript 0 means any given time. Similarly we may obtain the modified Friedmann equation of [4] in the form

$$\dot{R}_b^2 + \left(1 - \left(\frac{\rho + \sigma}{\sigma_c} \right)^2 \right) \frac{R_b^2}{\ell^2} - \frac{\mu}{R_b^2} = -k, \quad (18)$$

where $\sigma_c \equiv 3/(4\pi G\ell^2) = 6/(\kappa_5^2 \ell)$ is the critical tension for fine tuning. The cosmological constant in the brane then has the form

$$\frac{\Lambda_4}{3} = \frac{\kappa_5^4 \sigma^2}{36} - \frac{1}{\ell^2} = \frac{1}{\ell^2} \left(\frac{\sigma^2}{\sigma_c^2} - 1 \right).$$

4. Graviton emission and capture

4.1. Potential diagrams

In the case of a domain wall, there is no matter on the wall so we do not expect the escaping gravitons to be attracted back to the brane. A more realistic scenario would be to consider a brane with isotropically distributed matter. Analytically computing trajectories of the moving brane with matter is difficult so we make use of potential diagrams to study whether gravitational signals will return to the brane. In classical GR, potential diagrams have been found useful for qualitative analysis of orbits of particles and photons in Schwarzschild geometry. Potential diagrams has been used before to investigate Lorentz violations in $AdSS_5$ bulk background [20]. Here we extend that analysis to brane motion and graviton motion in AdS_5 and $AdSS_5$.

We start by rewriting equation (10) as

$$\left(\frac{dR}{d\lambda}\right)^2 + B^2(R) = b^2, \quad (19)$$

where

$$B^2(R) = \frac{f_k(R)}{R^2} \quad (20)$$

defines a kind of “effective potential of a graviton”, and

$$b^2 = \frac{E^2}{P^2} \quad (21)$$

defines the “impact parameter”.

The potential impact parameter $B(R)$ has the following useful interpretation: For a graviton to reach the point R , it must have a impact parameter b greater than or equal to $B(R)$, i.e.

$$b \geq B(R). \quad (22)$$

This is referred to as the *condition of accessibility* [16].

AdS_5 ($\mu = 0$): The effective potential (20) for the various values of k are

$$B^2(R) = \frac{1}{\ell^2} + \frac{1}{R^2}, \quad k = +1 \quad (23)$$

$$B^2(R) = \frac{1}{\ell^2}, \quad k = 0 \quad (24)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{1}{R^2}, \quad k = -1 \quad (25)$$

and are plotted in Figure 1. Clearly none of the potential curves have turning points, i.e if a graviton escape from the brane (domain wall or brane with matter) into an anti-de Sitter bulk, it will not return to the brane. For $k = -1$ we have a horizon at $R = \ell$.

$AdSS_5$ ($\mu \neq 0$): The effective potential (20) for the various values of k are

$$B^2(R) = \frac{1}{\ell^2} + \frac{1}{R^2} - \frac{\mu}{R^4}, \quad k = +1 \quad (26)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{\mu}{R^4}, \quad k = 0 \quad (27)$$

$$B^2(R) = \frac{1}{\ell^2} - \frac{1}{R^2} - \frac{\mu}{R^4}, \quad k = -1 \quad (28)$$




Figure 1. Potential diagram for AdS_5 ($\mu = 0$) bulk geometry.




Figure 2. Potential diagram for $\mu \neq 0$ and $k = +1$.

The case where $k = +1$ is plotted in Figure 2. Both the $k = 0$ and $k = -1$ cases look similar to $k = -1$ when $\mu = 0$ (see Figure 1), i.e they have no turning points. The horizons for all k 's are given by (3).

4.2. Condition of return

The turning point on Figure 2, is at

$$R_t = (2\mu)^{\frac{1}{2}} \quad (29)$$

which implies a maximum effective potential of

$$B^2(R_t) = \frac{1}{\ell^2} + \frac{1}{4\mu}.$$

Combining this with the condition of accessibility (22), we can summarize the qualitative features of the trajectories of gravitons emitted in the $AdSS_5$ bulk with $k = +1$ (see [16]) as follows:

When a graviton is emitted near the horizon R_h (i.e. at some point R_{em} , $R_h < R_{em} < R_t$), it will:

(i) escape to infinity when

$$b > \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}; \quad (30)$$

(ii) or return when

$$0 < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}. \quad (31)$$

Equation (31) in effect gives us the *condition of return* for a graviton emitted near the bulk horizon R_h . The lower limit arise from the fact that both E and P are positive constants, and hence $b > 0$. Clearly we can only expect returning gravitons at early times when $R_{em} < R_t$ when the effective potential is increasing. When a graviton is emitted at $R_{em} > R_t$ it will escape to infinity (since $B(R)$ is decreasing).

4.3. Conditions of escape

For a gravitational signal to escape from the brane, it must move faster than the brane with respect to an observer in the bulk. Thus the relative velocity of the gravitational signal v_g must be greater than the relative velocity of the brane v_b at the point of emission R_{em} . Therefore, we now seek to obtain the relative velocities of the gravitational signal and the brane.

We first obtain the relative velocity of the gravitational signal by dividing equation (8) into equation (10), and then taking the square root to find

$$v_g = \frac{dR_g}{dT} = f_k(R_g) \sqrt{1 - \frac{P^2 f_k(R_g)}{E^2 R_g^2}} = f_k(R_g) \sqrt{1 - \frac{B^2(R_g)}{b^2}}. \quad (32)$$

We see that this equation will hold if the condition of accessibility (22) is satisfied (where we made use of (20) and (21)). The velocity will be zero when $b = B(R_g)$, i.e. we will have a turning point for a gravitational signal.

Next we find the relative velocity of the brane by substituting equation (11) for dt into (16),

$$dR_b^2 = \left[\left(\frac{\rho + \sigma}{\sigma_c} \right)^2 \frac{R_b^2}{\ell^2} - f_k(R_b) \right] [f_k(R_b) dT^2 - f_k^{-1}(R_b) dR_b^2], \quad (33)$$

which on simplification yields

$$v_b = \frac{dR_b}{dT} = f_k(R_b) \sqrt{1 - \frac{\ell^2 f_k(R_b)}{\left(\frac{\rho + \sigma}{\sigma_c} \right)^2 R_b^2}}. \quad (34)$$

The equation for the relative velocity of the brane will only hold if

$$\hat{\beta} = \frac{1}{\ell} \left(\frac{\rho + \sigma}{\sigma_c} \right) \geq B(R_b). \quad (35)$$

In the limit of $R \rightarrow 0$ this imply that $\mu \geq 0$ and in the limit of $R \rightarrow \infty$ we must have $\sigma \geq \sigma_c$. We are therefore only considering positive dark radiation.

The ratio of the two relative velocities is then given by

$$\frac{v_g}{v_b} = \frac{f_k(R_g)}{f_k(R_b)} \cdot \sqrt{\frac{1 - \frac{P^2 f_k(R_g)}{E^2 R_g^2}}{1 - \frac{\ell^2 f_k(R_b)}{\left(\frac{\rho + \sigma}{\sigma_c}\right)^2 R_b^2}}}. \quad (36)$$

We seek to find the conditions of escape, and therefore we only consider the position of the brane and the energy density at the time of emission, i.e. $R_b = R_g = R_{em}$ and $\rho = \rho_{em}$ respectively. The graviton will have the same position as the brane at the bulk time of emission. A gravitational signal can be emitted if and only if $v_g > v_b$, that is when

$$1 - \frac{P^2 f_k(R_{em})}{E^2 R_{em}^2} > 1 - \frac{\ell^2 f_k(R_{em})}{\left(\frac{\rho_{em} + \sigma}{\sigma_c}\right)^2 R_{em}^2},$$

which after simplification reduces to

$$\frac{E}{P} > \frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) \quad (37)$$

or

$$b > \hat{\beta}_{em}. \quad (38)$$

This condition can be considered as the *condition of escape*, that is the condition which needs to be satisfied for a graviton to escape from the brane.

When $v_g = v_b$ we have the condition

$$\frac{E}{P} = \frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) \quad (39)$$

which implies that the graviton is stuck on the brane and therefore appears to be moving with the brane at the same relative velocity. In the case when $v_g < v_b$, the gravitational signal is lagging the brane. This is clearly not possible.

5. Summary and discussion

The only case which showed the possibility of emitting a graviton with it returning to the brane, is $k = +1$ when $\mu \neq 0$. The potential diagram for this case has a classical potential form (see [16] and Figure 2) which meant that if a graviton is emitted from the brane at R_{em} (where $R_h < R_{em} < R_t$) it would return if the condition (31) is satisfied. Comparing the velocities of the graviton and the brane we were able to determine that a graviton will be able to escape from the brane if the condition (37) is satisfied.

The condition for a graviton to escape from the brane at some early time (when $R_h < R_{em} < R_t$) and return, is then given by

$$\frac{1}{\ell} \left(\frac{\rho_{em} + \sigma}{\sigma_c} \right) < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}, \quad (40)$$

where $\mu > 0$ and $\sigma \geq \sigma_c$. This condition can also be expressed in terms of the 5-dimensional fundamental mass scale M_5 , by using the relationship $\kappa_5^2 = 8\pi G\ell = M_5^{-3}$, namely

$$\frac{\rho_{em} + \sigma}{6M_5^3} < b < \sqrt{\frac{1}{\ell^2} + \frac{1}{4\mu}}. \quad (41)$$

Since the impact parameter has no physical significance, we rather use the conditions above to find the relationship between parameters that do have physical significance. We find that the conditions of escape and return impose the following restrictions on the dark radiation μ (Schwarzschild-like mass)

$$\mu < \frac{\ell^2}{4} \left[\left(\frac{\rho_{em} + \sigma}{\sigma_c} \right)^2 - 1 \right]^{-1}, \quad (42)$$

for all $\rho_{em} + \sigma > \sigma_c$. This in turn tells us that we may only have returning gravitons before the brane reaches the point R_t , where using (29) we have the condition

$$R_t < \frac{\ell}{\sqrt{2}} \left[\left(\frac{\rho_{em} + \sigma}{\sigma_c} \right)^2 - 1 \right]^{-\frac{1}{2}}. \quad (43)$$

We can realistically only expect emitted gravitons to return to the brane at very early times ($\rho_{em} + \sigma \gg \sigma_c$) when $R_{em} \rightarrow R_h$ and thus $B(R_{em}) \rightarrow 0$. At late times ($\rho \rightarrow 0$) we are essentially looking at a domain wall moving in an AdS_5 space where gravitons do not return to the brane (see appendix). In the limit $\rho_{em} + \sigma \gg \sigma_c$ the restrictions above reduce to

$$\mu < \frac{1}{(\rho_{em} + \sigma)^2} \quad (44)$$

and hence

$$R_t < \frac{1}{\rho_{em} + \sigma}. \quad (45)$$

The upper limit for the dark radiation μ obtained here still require further investigation, especially in the light of the nucleosynthesis limits [21].

We see that there exist a small window of opportunity, before the brane reaches R_t , for gravitons to be emitted from the brane and then return to it. After the brane reaches R_t , all gravitons emitted never return to the brane. An alternative might be to look at a bulk with some sort of field (dilaton etc.) which forces the path back to the brane when it has gone beyond R_t . Another possibility is to have a black hole on the brane that will give us the required curvature. Black holes on the brane have been consider by a number of authors (see [22–26]) most notably in the context of causality, by [27].

Another issue that needs investigation, is that of conservation of energy for a brane that is radiating these graviton. Clearly the form (15) does not take into consideration the gravitons that are emitted. Radiating braneworlds have been considered in [28] and [29], which might show more promising results. The mechanisms that could produce gravitons with sufficient energy to leave the brane has also as yet not been explored properly.

In conclusion, there are two important aspect of escaping gravitational signals. In §4 we clearly saw that the condition of return does not depend on the type of brane that we choose, but rather on the type of bulk (i.e. on its 5-D spacetime). We

also discovered that the condition of escape depends on the type of brane we choose and is independent of the bulk space time. We note that no matter which values of μ (AdS_5) and charge Q (Reissner-Nordström) we choose, the escape condition still stays the same. The condition of escape is also independent of the curvature values k .

We have thus shown that there are certain conditions which first needs to be satisfied before one may assume that a signal can escape and then return to the brane. Two important conditions need to be satisfied, firstly we need a condition under which a graviton can escape from the brane (or be captured by the brane), and secondly the conditions which allow it to return to the brane. Furthermore, these two conditions together restrict gravitons to escaping and returning to the brane before the brane reaches a point R_t in the bulk. If all of these conditions are not satisfied one can not expect to have causally connected points through the bulk.

Acknowledgments

This research was supported by the National Research Foundation (South Africa). J.L. also acknowledges the postgraduate assistantship from the University of South Africa (UNISA).

Appendix A. Domain wall motion in AdS_5

In this section we give the results for the special case where we are considering a empty brane (domain wall) moving in an AdS_5 bulk.

Appendix A.1. Null geodesics solutions for AdS_5

When the bulk is anti-de Sitter AdS_5 , i.e $\mu = 0$, one can readily obtain the solutions of the null geodesic equations (8)-(10). For $k = 0$ the solutions are

$$R_g = \frac{\mathcal{B}_g}{\mathcal{A}_g - T}, \quad \alpha > 0, \quad (\text{A.1})$$

and

$$R_g = \frac{\mathcal{B}_g}{T - \mathcal{A}_g}, \quad \alpha < 0, \quad (\text{A.2})$$

where $\alpha^2 = \frac{E^2}{P^2} - \frac{1}{\ell^2}$ and

$$\mathcal{A}_g = T_0 \pm \frac{E\ell^2}{\alpha P R_0}, \quad \mathcal{B}_g = \frac{E\ell^2}{\alpha P}.$$

The + sign in $\mathcal{A}_g$ holds when $\alpha > 0$, the - sign when $\alpha < 0$.

In the case where $k = -1$, the solutions are

$$R_g^2 = \frac{E^2\ell^2}{\alpha^2 P^2} \coth^2 \left(\frac{\mathcal{C}_g - T}{\ell} \right) - \frac{1}{\alpha^2}, \quad R > \ell, \quad (\text{A.3})$$

and

$$R_g^2 = \frac{E^2\ell^2}{\alpha^2 P^2} \tanh^2 \left(\frac{\mathcal{D}_g - T}{\ell} \right) - \frac{1}{\alpha^2}, \quad R < \ell, \quad (\text{A.4})$$

where the initial conditions are

$$\mathcal{C}_g = T_0 + \ell \coth^{-1} \frac{P\sqrt{\alpha^2 R_0^2 + 1}}{E\ell},$$

and

$$\mathcal{D}_g = T_0 + \ell \tanh^{-1} \frac{P\sqrt{\alpha^2 R_0^2 + 1}}{E\ell}.$$

Similarly for $k = +1$ the solutions are

$$R_g^2 = \frac{E^2 \ell^2}{\alpha^2 P^2} \tan^2 \left(\frac{T - \mathcal{F}_g}{\ell} \right) + \frac{1}{\alpha^2}, \quad (\text{A.5})$$

where

$$\mathcal{F}_g = T_0 - \ell \tan^{-1} \frac{P\sqrt{\alpha^2 R_0^2 - 1}}{E\ell}.$$

The solutions to the geodesic equation for the radial coordinate r in terms of the global bulk time T , for all $k = 0, -1$ and $+1$, are given by

$$r = r_0 + \frac{P}{E\ell^2} (T - T_0). \quad (\text{A.6})$$

Appendix A.2. Trajectories for a moving domain wall

When one considers a domain wall (i.e. $\rho = 0$) moving in an anti-de Sitter bulk ($\mu = 0$), the modified Friedmann equation (18) takes the form [8]

$$\dot{R}_b^2 + \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \right) \frac{R_b^2}{\ell^2} = -k. \quad (\text{A.7})$$

The solutions of these, in terms of the proper time on the brane t , are given in Kraus [8]. We are however interested in the brane's trajectory in the bulk. We therefore transform Kraus' equations by making use of (12). The trajectories are

$k = 0, \sigma = \sigma_c$: $R_b = \text{constant}$

$k = 0, \sigma > \sigma_c$:

solution 1 (expanding brane):

$$R_b = \frac{1}{H} \left(\frac{\sigma}{\sigma_c} \right) \frac{\ell}{\mathcal{A}_b - T}, \text{ where } \mathcal{A}_b = T_0 + \frac{\sigma}{\sigma_c} \frac{\ell}{R_0 H}, \quad (\text{A.8})$$

and where $H = \frac{1}{\ell} \sqrt{\left(\frac{\sigma}{\sigma_c} \right)^2 - 1}$.

solution 2 (contracting brane):

$$R_b = \frac{1}{H} \left(\frac{\sigma}{\sigma_c} \right) \frac{\ell}{T - \mathcal{A}_b}, \text{ where } \mathcal{A}_b = T_0 - \frac{\sigma}{\sigma_c} \frac{\ell}{R_0 H}. \quad (\text{A.9})$$

$k = -1, \sigma = \sigma_c$:

$$R_b^2 = \ell^2 - (\ell^2 - R_0^2) e^{\frac{2(T-T_0)}{\ell}}, \quad R < \ell, \quad (\text{A.10})$$

and

$$R_b^2 = (R_0^2 - \ell^2) e^{\frac{2(T-T_0)}{\ell}} + \ell^2, \quad R > \ell. \quad (\text{A.11})$$

$k = -1, \sigma > \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \coth^2 \left(\frac{\mathcal{C}_b - T}{\ell} \right) - 1 \right), \quad R > \ell, \quad (\text{A.12})$$

and

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \tanh^2 \left(\frac{\mathcal{D}_b - T}{\ell} \right) - 1 \right), \quad R < \ell, \quad (\text{A.13})$$

where

$$\mathcal{C}_b = T_0 + \ell \coth^{-1} \left(\frac{\sqrt{H^2 R_0^2 + 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right),$$

and

$$\mathcal{D}_b = T_0 + \ell \tanh^{-1} \left(\frac{\sqrt{H^2 R_0^2 + 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$

$k = -1, \sigma < \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \coth^2 \left(\frac{T - \mathcal{C}_{b2}}{\ell} \right) \right), \quad R < \ell, \quad (\text{A.14})$$

and

$$R_b^2 = \frac{1}{H^2} \left(1 - \left(\frac{\sigma}{\sigma_c} \right)^2 \tanh^2 \left(\frac{T - \mathcal{D}_{b2}}{\ell} \right) \right), \quad R > \ell, \quad (\text{A.15})$$

where

$$\mathcal{C}_{b2} = T_0 - \ell \coth^{-1} \left(\frac{\sqrt{1 - H^2 R_0^2}}{\left(\frac{\sigma}{\sigma_c} \right)} \right),$$

and

$$\mathcal{D}_{b2} = T_0 - \ell \tanh^{-1} \left(\frac{\sqrt{1 - H^2 R_0^2}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$

$k = +1, \sigma > \sigma_c$:

$$R_b^2 = \frac{1}{H^2} \left(\left(\frac{\sigma}{\sigma_c} \right)^2 \tan^2 \left(\frac{T - \mathcal{F}_b}{\ell} \right) + 1 \right), \quad (\text{A.16})$$

where

$$\mathcal{F}_b = T_0 - \ell \tan^{-1} \left(\frac{\sqrt{H^2 R_0^2 - 1}}{\left(\frac{\sigma}{\sigma_c} \right)} \right).$$




Figure A1. Expanding brane: R versus T for $k = 0$, $\sigma > \sigma_c$ and $\alpha > 0$.

Appendix A.3. Analysis

We now have the trajectories for both the empty brane (domain wall) and an gravitational signal in AdS_5 . We plot them on the same set of axes, with the same initial conditions so as to see the behaviour of the gravitational signal after it has left the brane.

In order for us to compare the trajectories we set

$$\omega = \frac{E\ell}{P} \quad (\text{A.17})$$

in the null geodesic equations, and set

$$\beta = \frac{\sigma}{\sigma_c} \quad (\text{A.18})$$

in the brane's equations of motion. We assume that $\ell \sim 10^{-4}m$.

In Figure A1 we start with the $k = 0$ case for an expanding universe (solution 1) with $\sigma > \sigma_c$. We set $\beta = 2$ and $R_0 = 1$ at $T_0 = 0$, in (A.8) for the brane equation of motion. For the gravitational signals path (null geodesic), we consider $\omega = 4$ and $\omega = 1.5$ in (A.1) with the same initial conditions. In Figure A2 we plot the results for a contracting universe (solution 2). The two cases for $k = -1$ are plotted in Figure A3 and Figure A4 and the case when $k = 1$ is plotted in Figure A5 (all with $\sigma > \sigma_c$).

Appendix A.3.1. $\omega = \beta$ When we set $\omega = \beta$ we find that the gravitational signal follow the same path as the brane for all cases of k when $\sigma > \sigma_c$. We can interpreted this condition as the graviton being confined to the brane and therefore having the same coordinate in the bulk. We can restate this condition in terms of the impact parameter

$$b = \frac{E}{P} = \frac{1}{\ell} \left(\frac{\sigma}{\sigma_c} \right) = \frac{\sigma}{6M_5^3}. \quad (\text{A.19})$$




Figure A2. Contracting brane: R versus T for $k = 0$, $\sigma > \sigma_c$ and $\alpha < 0$.




Figure A3. R versus T for $k = -1$ and $\sigma > \sigma_c$ and $R > \ell$




Figure A4. R versus T for $k = -1$, $\sigma > \sigma_c$ and $R < \ell$.




Figure A5. R versus T for $k = 1$ and $\sigma > \sigma_c$

Appendix A.3.2. $\omega > \beta$ When $\omega > \beta$ the geodesic equation is not the same as the equation of motion. Thus a graviton can be emitted along the extra-dimensional geodesic, i.e. can be emitted from the brane. This can therefore be seen as the condition for a graviton to be emitted from a domain wall with tension σ into a AdS_5 space. We can express this condition in the form

$$b = \frac{E}{P} > \frac{1}{\ell} \left(\frac{\sigma}{\sigma_c} \right). \quad (\text{A.20})$$

This condition amounts to the gravitational signal having a greater velocity than the brane and may serve as a late time approximation of (37) since at late times the AdS_5 becomes AdS_5 and the brane a domain wall. We can therefore see from Figure A5 that no gravitational signals return to the brane at late times.

Appendix A.3.3. $\omega < \beta$ In the $k = 0$ and $k = -1$ cases we see that the possibility of $\omega < \beta$ also gives paths that are different to the brane's path. However, in these cases the brane is moving faster than the gravitational signal. Clearly these cases are not valid. The initial conditions for the $k = 1$, exclude the possibility of $\omega < \beta$.

References

- [1] Chung D and Freese K, *Phys. Rev.* **D62**, 063513 (2000).
- [2] Ishihara H, *Phys. Rev. Lett.* **86**, 381 (2001).
- [3] Binétruy P, Deffayet C and Langlois D, *Nuc. Phys.* **B565**, 269 (2000).
- [4] Binétruy P, Deffayet C, Ellwanger U and Langlois D, *Phys. Lett.* **B477**, 285 (2000).
- [5] Youm D, *Mod. Phys. Lett.* **A16**, 2371 (2001).
- [6] Davis A, Rhodes C and Vernon I, *JHEP* **0111**, 015 (2001).
- [7] Abdalla E, Cuadros-Melgar B, Feng S and Wang B, *Phys. Rev.* **D65**, 083512 (2002).
- [8] Kraus P, *JHEP* **9912**, 011 (1999).
- [9] Bowcock P, Charmousis C and Gregory R, *Class. Quant. Grav.* **17**, 4745 (2000).
- [10] Chamblin A and Reall H S, *Nuc. Phys.* **B562**, 133 (1999).
- [11] Caldwell R and Langlois D, *Phys. Lett.* **B511**, 129 (2001).
- [12] Abdalla E, Casali A G and Cuadros-Melgar B, *Int. J. Theor. Phys.* **43**, 801 (2004).
- [13] Abdalla E, Casali A G and Cuadros-Melgar B, *Nuc. Phys.* **B644**, 201 (2002).
- [14] Abdalla E and Cuadros-Melgar B, *Phys. Rev.* **D67**, 084012 (2003).
- [15] Cuadros-Melgar B, *Class. Quant. Grav.* **21**, 2669 (2004).
- [16] Misner C W, Thorne K S and Wheeler J A, *Gravitation* (Freeman, San Francisco 1973).
- [17] Israel W, *Nuovo Cimento* **B44**, 1 (1966), erratum: **B48**, 463 (1967).
- [18] Brax P, Langlois D and Rodríguez-Martínez M, *Phys. Rev.* **D67**, 104022 (2003).
- [19] Brax P and van de Bruck C, *Class. Quant. Grav.* **20**, R201 (2003) [arXiv: hep-th/0303095].
- [20] Csáki C, Erlich J and Grojean C, *Nuc. Phys.* **B604**, 312 (2001).
- [21] Ichiki K, Yahiro M, Kajino T, Orito M and Mathews G J, *Phys. Rev.* **D66**, 043521 (2002).
- [22] Chamblin A, Hawking S W and Reall H S, *Phys. Rev.* **D61**, 065007 (2000).
- [23] Dadhich N, Maartens R, Papadopoulos P and Rezanian V, *Phys. Lett.* **B487**, 1 (2000).
- [24] Emparan R, Horowitz G T and Myers R C, *JHEP* **0001**, 007 (2000).
- [25] Chamblin A, Reall H.S, Shinkau H. and Shiromizu T., *Phys. Rev.* **D63**, 064015 (2001).
- [26] Pravda V and Pravdova A, *Class. Quant. Grav.* **18**, 1205 (2001).
- [27] Chamblin A, *Class. Quant. Grav.* **18**, L17 (2001).
- [28] Langlois D, Sorbo L and Rodríguez-Martínez M, *Phys. Rev. Lett.* **89**, 171301 (2002).
- [29] Leeper E, Maartens R and Sopuerta C F, *Class. Quant. Grav.* **21**, 1125 (2004).